Neil L. Duplantier, M.D Bone and Joint Clinic www.boneandjointclinic.com



## Rehabilitation Protocol: **Distal Femoral Osteotomy (DFO)**

	Weight Bearing	Brace Use	Range of Motion	Therapeutic Elements
PHASE I 0 -2 weeks 2-6 weeks	Touch-down weight bearing (TDWB) with crutches.	Locked in extension at all times including sleep. Remove for exercises and possible CPM.  Unlocked.	Advance as tolerated. Goal: at least 90 degs by 4wks, 120 degs by 6wks.  PROM: 90 degs flexion at each therapy session.	Quad sets, hamstring isometrics, co-contractions, ankle pumps. Regain full extension. Modalities: heat before, ice after. Quad stim.
PHASE II 6-8 weeks	As tolerated (WBAT), wean off crutches.	Discontinue brace if patient has controlled SLR without extension lag.	Progress to full ROM.	Start SLR out of brace. Progressive closed chain exercise. Bicycle without resistance.
8-12 weeks	Full weight (FWB) with normal gait.	None.	Full A/PROM.	Advance closed chain exercise. Leg press 0-60 degs, Short are wall squats. Balance exercises.
PHASE III 12 -16 weeks	FWB.	None.	Full A/PROM.	Begin open chain strengthening. Swimming and Treadmill walking for exercise. Progress to Elliptical (EFX).
PHASE IV 4-6 months	FWB.	None.	Full.	Start progressive running program. Sports specific drills.  Continue strength training.  Consider return to high impact sports.

## **LOCATIONS**

2600 Belle Chasse Hwy Gretna, LA, 70056

Tel: 504-391-7670 Fax: 504-378-9439 4633 Wichers Drive Marrero, LA, 70072 Tel: 504-347-5421

Fax: 504-340-5171