



Rehabilitation Protocol: **Osteochondral Allograft Transplantation (Femoral Condyle)**

	Weight Bearing	Brace Use	Range of Motion	Therapeutic Elements
PHASE I 0 - 2 weeks	Non-weight bearing (NWB) with crutches.	Locked in extension. Remove for CPM and exercises.	CPM: 6-8 hrs / day. Start -5 to 45 degrees. Increase 5-10 degs / day as tolerated.	Quad sets, SLR, hamstring isometrics, ankle pumps. Regain full extension. Hamstring stretching. Patella mobs.
2 - 6 weeks	Touch-down (TDWB) with crutches.	Brace progressively opened as quad control returns.	P/AROM as tol. Goal: at least 90 degs by 4wks, 120 degs by 6wks.	
PHASE II 6 - 8 weeks	As tolerated (WBAT) with crutches	Discontinue brace if patient has controlled SLR without extension lag.	Full A/PROM.	Begin progressive closed chain exercise. Bicycle.
8 - 12 weeks	Full weight bearing (FWB). Wean off crutches.	None.	Full A/PROM.	Gait training. Closed chain exercise. Balance exercises.
PHASE III 3 - 6 months	FWB with normal gait.	None.	Full A/PROM.	Advance Begin open chain strengthening. Swimming and Treadmill walking for exercise. Progress to Elliptical (EFX).
PHASE IV 6-12 months	FWB.	None.	Full.	Continue strength training. May begin a progressive running and agility program. Consider return to high impact sports.

LOCATIONS

2600 Belle Chasse Hwy
Gretna, LA, 70056
Tel: 504-391-7670
Fax: 504-378-9439

4633 Wichers Drive
Marrero, LA, 70072
Tel: 504-347-5421
Fax: 504-340-5171