Neil L. Duplantier, M.D **Bone and Joint Clinic** www.boneandjointclinic.com



Rehabilitation Protocol: **Arthroscopic Partial Meniscectomy / Debridement**

	Weight Bearing	Range of Motion	Therapeutic Elements
PHASE I 0 – 2 weeks	Weight bearing as tolerated (WBAT). Crutches used for 24-48 hrs. Advance to full weight bearing as tolerated.	No restrictions on motion. Focus on immediate full extension. Advance to full flexion as tolerated.	Heel slides, Quad and hamstring sets, Patella mobilizations, Straight leg raises, Prone hangs. ICE 3x/day x 20 minutes (Do not place directly on exposed skin) Modalities as needed.
PHASE II 2 – 4 weeks	Full weight bearing	Full ROM	Stationary bike. Wall sits and lunges. Balancing exercises.
PHASE III 4 - 6 weeks	Full weight bearing	Full ROM	Advance closed chain strengthening, leg press, leg curls. Plyometric and proprioceptive training.

LOCATIONS

2600 Belle Chasse Hwy Gretna, LA, 70056 Tel: 504-391-7670

Fax: 504-378-9439

4633 Wichers Drive Marrero, LA, 70072

Tel: 504-347-5421 Fax: 504-340-5171