



Rehabilitation Protocol: Shoulder Replacement (Total and Hemiarthroplasty)

	RANGE OF MOTION*	IMMOBILIZER	THERAPEUTIC EXERCISE
PHASE I 0 - 6 weeks	Passive to Active-assisted to Active as tolerated within limits: FF >= 120 degs ER= max 40 degs at side ABD= max 75 degs without rotation No active IR or EXT	Worn at all times except for hygiene and therapeutic exercise for 4-6 wks.	Elbow/wrist/hand ROM, grip strength. Pendulums. Isometric scapular stabilizers exercises. Canes and Pulleys. Modalities: heat before, ice after.
PHASE II 6 - 12 weeks	Continue PROM stretch at end ranges. Add A/AAROM as tolerated for IR and extension.	Discontinue sling at 6wks.	Begin light resisted ER/FF/ABD. Isometrics and therabands. (concentric motions only). No resisted IR/EXT or scapular retractions.
PHASE III 3 - 6 months	Progress to full active motion. No restrictions.	None.	Adv strengthening. Add IR/EXT and scapular retraction strengthening. Begin eccentric motions, plyometrics, closed chain exercises.
PHASE IV 6 - 12 months	Full painless motion.	None.	Begin functional sport/work specifics. Return to previous activity level.

LOCATIONS

**2600 Belle Chasse Hwy
 Gretna, LA, 70056
 Tel: 504-391-7670
 Fax: 504-378-9439**

**4633 Wichers Drive
 Marrero, LA, 70072
 Tel: 504-347-5421
 Fax: 504-340-5171**